



ELATUS[®] Co-Pack is a combination of separately registered products comprised of ELATUS A and ELATUS B, A Solatenol[®] Brand Fungicide.

ELATUS[®], Solatenol[®] and the Syngenta wordmark are trademarks of a Syngenta Group Company.

GROUP

11

FUNGICIDE

ELATUS® A Fungicide

SUSPENSION

COMMERCIAL

For Use in Controlling Diseases in Labelled Crops.

ACTIVE INGREDIENT:

Azoxystrobin..... 250 g/L

Contains 1,2-benzisothiazolin-3-one at 0.031% OR 1,2-benzisothiazolin-3-one at 0.039% plus 2-bromo-2-nitropropane-1,3-diol at 0.03% OR 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00089% and 2-methyl-4-isothiazolin-3-one at 0.00030% plus 2-bromo-2-nitropropane-1,3-diol at 0.003% as preservatives.

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

REGISTRATION NO: **31973**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 L – 1000 L**

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

Label

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

3.0 TOXICOLOGICAL INFORMATION

No specific symptoms of poisoning are known for this product. If ingested, nausea, vomiting, diarrhea and abdominal pain may occur. Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

May irritate eyes. Avoid contact with eyes, skin and clothing. Avoid breathing dust or spray mist. Wash with soap and water after handling, and before eating, drinking or smoking. Wash contaminated clothing, separately from household laundry, before re-use. Do not wear contaminated shoes.

RESTRICTED ENTRY INTERVAL (REI): Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours following application.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt and long pants when mixing, loading and applying and during clean-up and repair activities. Wear chemical resistant gloves during mixing and loading.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Azoxystrobin is persistent and will carryover. It is recommended that this product not be used in areas treated with Azoxystrobin during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

7.0 STORAGE

Keep in original container, tightly closed, during storage. Store this product away from food or feed.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

ELATUS® is a trademark of a Syngenta Group Company.

GROUP

11

FUNGICIDE

ELATUS® A Fungicide

SUSPENSION

COMMERCIAL

For Use in Controlling Diseases in Labelled Crops.

ACTIVE INGREDIENT:

Azoxystrobin..... 250 g/L

Contains 1,2-benzisothiazolin-3-one at 0.031% OR 1,2-benzisothiazolin-3-one at 0.039% plus 2-bromo-2-nitropropane-1,3-diol at 0.03% OR 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00089% and 2-methyl-4-isothiazolin-3-one at 0.00030% plus 2-bromo-2-nitropropane-1,3-diol at 0.003% as preservatives.

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

REGISTRATION NO: **31973**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

Pamphlet

Table of Contents

	<u>Section Number</u>
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE)	5.0
ENVIRONMENTAL PRECAUTIONS	6.0
STORAGE	7.0
DISPOSAL	8.0
PRODUCT INFORMATION	9.0
DIRECTIONS FOR USE	10.0
General Information	10.1
Ground Application	10.2
Mixing Instructions	10.2.1
Spraying Instructions	10.2.2
Equipment Clean-Up	10.2.3
Aerial Application	10.3
Use Precautions	10.3.1
Operator Precautions	10.3.2
Mixing Instructions	10.3.3
Spraying Instructions	10.3.4
Equipment Clean-Up	10.3.5
Product Specific Precautions	10.3.6
Spray Buffer Zones	10.4
Rotational Crop Restrictions	10.5
CROP USE DIRECTIONS	11.0
Potatoes – In Furrow Application	11.1
Legume Vegetables (Crop Group 6) Including Soybeans	11.2
MINOR USES	12.0
Sugar Beets	12.1
RESISTANCE MANAGEMENT RECOMMENDATIONS	13.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

3.0 TOXICOLOGICAL INFORMATION

No specific symptoms of poisoning are known for this product. If ingested, nausea, vomiting, diarrhea and abdominal pain may occur. Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

May irritate eyes. Avoid contact with eyes, skin and clothing. Avoid breathing dust or spray mist. Wash with soap and water after handling, and before eating, drinking or smoking. Wash contaminated clothing, separately from household laundry, before re-use. Do not wear contaminated shoes.

RESTRICTED ENTRY INTERVAL (REI): Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours following application.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt and long pants when mixing, loading and applying and during clean-up and repair activities. Wear chemical resistant gloves during mixing and loading.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Azoxystrobin is persistent and will carryover. It is recommended that this product not be used in areas treated with Azoxystrobin during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

7.0 STORAGE

Keep in original container, tightly closed, during storage. Store this product away from food or feed.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

9.0 PRODUCT INFORMATION

ELATUS® A Fungicide is a broad spectrum, preventative fungicide with systemic properties recommended for the control of plant diseases of legume vegetables (Crop Group 6), including soybeans.

ELATUS A Fungicide is to be applied as a foliar spray. Optimum disease control/suppression will be obtained by protective treatments prior to disease establishment. Refer to the specific use directions for each crop for detailed instructions on rates, application timing and technique.

Do not apply ELATUS A Fungicide through irrigation equipment.

10.0 DIRECTIONS FOR USE**10.1 General Information**

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of waste.

10.2 Ground Application

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply spray droplets which are smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

10.2.1 Mixing Instructions

1. Ensure that the sprayer interior is clean, then fill the spray tank with $\frac{1}{2}$ the required amount of water and engage gentle agitation. Good agitation is indicated by a rippling or rolling action on the surface of the water.
2. Add any WG or DF formulation mix partners and agitate to ensure complete mixing.
3. Add ELATUS A (SC) and agitate to ensure complete mixing.
4. Add any additional SC formulation mix partners and agitate to ensure complete mixing.
5. Add any EC formulation mix partners and agitate to ensure complete mixing.
6. Fill the tank to $\frac{3}{4}$ the required amount of water.
7. Add any solution (SN or SL) formulation mix partners and agitate to ensure complete mixing.
8. Finish filling the sprayer with water, maintaining good agitation.
9. After any break in spraying operations, agitate thoroughly before spraying again.
10. Spray the pesticide suspension the same day as mixing.
11. Do not mix, load or clean spray equipment where there is a potential to contaminate wells or aquatic systems.

When using chemical handling equipment to fill the sprayer, the following additional recommendations apply:

- WG and DF formulations are preferentially batch mixed.
- SC, SN, and SL formulations may be inducted or batch mixed.
- EC formulations are preferentially batch mixed.

10.2.2 Spraying Instructions

1. Water Volume: Specific to crop and disease. Consult the following tabulated instructions for use.
2. Spray Nozzles: 80° or 110° drift reducing flat fan (e.g. those with a pre-orifice or turbulence chamber) or air induction nozzles are recommended. Use 50 mesh nozzle screens. Do not use flood type nozzles, controlled droplet application equipment, spray foils or hollow cone nozzles.
3. Pressure: As recommended by the nozzle manufacturer to achieve ASAE medium sized droplets.
4. Apply at uniform speed and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to avoid potential crop injury from over application.

10.2.3 Equipment Clean-Up

Before Spraying:

- Prior to using ELATUS A fungicide, ensure that the spray tank, lines and filter are thoroughly clean.

After Spraying:

- Thoroughly clean application equipment immediately after spraying. Do not allow ELATUS A residue to dry within the spray tank.
- When using tank mixes, consult the tank-mix partner label for additional cleanup instructions.
- The following recommendations are provided:
 1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **Do not** clean the sprayer near desirable vegetation, wells or other water sources.

2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial regulations.

10.3 Aerial Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply spray droplets which are smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions, and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. When no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking technology. GPS based marking is recommended.

10.3.1 Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

10.3.2 Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

10.3.3 Mixing Instructions

Mixing this product directly in the aircraft hopper **IS NOT** recommended. The use of chemical handling or managing equipment to load the hopper **IS** recommended. This product **MAY BE** inducted into a hopper which is prefilled with water or when the product and water are mixed prior to

entering the hopper. This product **MAY BE** batch mixed and pumped into the hopper. In all cases the chemical handling equipment and hopper interior must be clean prior to use.

NOTE: WG and DF formulations are preferentially batch mixed.

NOTE: SC, SN, and SL formulations may be inducted or batch mixed.

NOTE: EC formulations are preferentially batch mixed.

It is **NOT** recommended to combine solid (WG or DF) formulations with liquid tank mix partners within a single batch. Batch mix WG or DF formulations first, pump into the hopper, and then add liquid tank mix partners by induction or as an additional batch mix. When tank mixing multiple products, follow the mixing order outlined below:

1. Pump water into the hopper to at least $\frac{1}{4}$ to $\frac{1}{2}$ of the desired spray volume. Engage hopper circulation, if possible.
2. Thoroughly batch mix any WG or DF formulation mix partners and agitate to ensure complete mixing. Pump into the hopper.
3. Induct or thoroughly batch mix ELATUS A (SC) and any additional SC formulations.
4. Thoroughly batch mix any EC formulation mix partners.
5. Induct or thoroughly batch mix any solution (SN or SL) formulation mix partners.
6. Pump batch mixed SC, EC, and/or SN/SL products into the hopper.
7. Finish filling the hopper with water.
8. If it was not possible to engage hopper agitation in Step 1, do so as soon as possible once airborne.
9. Spray the pesticide suspension the same day as mixing.
10. Do not mix, load or clean equipment where there is a potential to contaminate wells or aquatic systems.

10.3.4 Spraying Instructions

1. Water Volume: Apply in a minimum spray volume of 45 L/ha OR the volume given in the crop and pest specific instructions tabulated below, whichever is LARGER.
2. Spray Nozzles: Use only ASAE medium or coarse nozzles rated as delivering droplets of volume median diameter of 300 microns or greater.
3. Pressure: As recommended by the nozzle manufacturer to achieve ASAE medium sized droplets.
4. Ensure hopper agitation is engaged whenever possible during flight.

10.3.5 Equipment Clean-Up

Before Spraying:

- Prior to using ELATUS A fungicide, ensure that the hopper, chemical handling equipment, lines and filter are thoroughly cleaned.

After Spraying:

- Thoroughly clean application equipment immediately after spraying. Do not allow ELATUS A residue to dry within application equipment.
- When using tank mixes, consult the tank-mix partner label for additional cleanup instructions.
- The following recommendations are provided:
 1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **Do not** clean application equipment near desirable vegetation, wells or other water sources.

2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial regulations.

10.3.6 Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-87-SYNGENTA (1-877-964-3682) or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum of 45 litres of water per hectare.

ELATUS A Fungicide is extremely phytotoxic to certain apple varieties. Avoid spray drift. Extreme care must be taken to prevent injury to apple trees (and apple fruit). DO NOT spray ELATUS A where spray drift may reach apple trees.

Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Follow recommendations provided by local disease monitoring services or provincial spray calendars regarding the appropriate application timing for protectant fungicides in your area. Cultural practices such as canopy management and removal of overwintered plant debris should be integrated with the use of fungicides to reduce disease incidence. Use the higher rate, and the shorter application interval under conditions of heavy infection pressure, on highly susceptible varieties, or when environmental conditions are favourable for disease development.

ELATUS A Fungicide treatments should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, plant residue management, crop rotation, and proper timing and placement of fertilizer and irrigation. Fungicide applications should begin prior to disease infection and continue throughout the season following a resistance management strategy (see **Resistance-Management Recommendations**).

10.4 Spray Buffer Zones

Use of the following spray methods or equipment DO NOT require a spray buffer zone: hand-held or backpack sprayer and spot treatment, inter-row hooded sprayer, soil drench and soil incorporation.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:			
		Freshwater Habitats of Depths:		Estuarine/Marine Habitats of Depths:	
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field sprayer	Soybeans, Legumes, Potato, Sugarbeets	1	1	1	1
Aerial	Soybeans, Legumes	5	1	1	1

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticide portion of the Canada.ca web site.

10.5 Rotational Crop Restrictions

Please see the following table for the crop rotational restrictions:

Rotational Crops	Planting Time From Last Application of Azoxystrobin-containing Products
All crops with Azoxystrobin registered uses	0 days
All other crops Intended for Food and Feed	30 days

11.0 CROP USE DIRECTIONS

11.1 Potatoes - In-Furrow Application

POTATOES – IN-FURROW APPLICATION	
DISEASE CONTROLLED	Soilborne Diseases: Silver Scurf (<i>Helminthosporium solani</i>) Rhizoctonia Stem Canker (<i>Rhizoctonia sp.</i>) Rhizoctonia Stolon Canker (<i>Rhizoctonia sp.</i>) Black Scurf (<i>Rhizoctonia solani</i>)
PRODUCT RATE	4-6 mL product/100 m row
NOTES	Apply once as an in furrow spray in 50 to 140 L per hectare of water at planting. Mount the spray nozzle so the spray is directed into the furrow as a 15-20 cm band just before the seed is covered.
Restrictions:	
1. DO NOT apply by air.	

11.2 Legume Vegetables (Crop Group 6) Including Soybeans

DISEASE	Asian (Soybean) Rust (<i>Phakopsora pachyrhizi</i>)
CROPS	<p>Crop Subgroup 6A: Edible-podded legume vegetables - Any succulent cultivar of edible podded bean (<i>Phaseolus</i> spp.) and any succulent cultivar of edible-podded pea (<i>Pisum</i> spp.). Bean (<i>Phaseolus</i> spp.) (includes runner bean, snap bean, wax bean); bean (<i>Vigna</i> spp.) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean); jack bean; pea (<i>Pisum</i> spp.) (includes dwarf pea, edible-podded pea, snow pea, sugar snap pea); pigeon pea; soybean (immature seed); and sword bean.</p> <p>Crop Subgroup 6B: Succulent shelled pea and bean - Any succulent shelled cultivar of bean (<i>Phaseolus</i> spp.) and garden pea (<i>Pisum</i> spp.); bean (<i>Phaseolus</i> spp.) (includes lima bean); broad bean (succulent fava bean); bean (<i>Vigna</i> spp.) (includes black-eyed pea, cowpea, southern pea); pea (<i>Pisum</i> spp.) (includes English pea, garden pea, green pea); and pigeon pea.</p> <p>Crop Subgroup 6C: Dried shelled pea and bean (except soybean) - Any dried cultivar of bean (<i>Phaseolus</i> spp.); and dried cultivar of pea (<i>Pisum</i> spp.); dried cultivars of bean (<i>Lupinus</i> spp.) (includes grain lupin, sweet lupin, white lupin, and sweet white lupin); bean (<i>Vigna</i> spp.) (includes adzuki beans, mung bean, rice bean, southern bean, urd bean); bean (<i>Phaseolus</i> spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); broad bean (dry, fava bean); chickpea; guar; lablab bean; lentil; pea (<i>Pisum</i> spp.) (includes field pea); and pigeon pea.</p> <p>Soybeans</p>
PRODUCT RATE (mL/ha)	500
APPLICATION TIMING	Make first application at the R1 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.
DISEASE	Powdery Mildew (<i>Microsphaera diffusa</i> , <i>Erysiphe pisi</i> , <i>E. polygoni</i>)
CROPS	Soybeans, and field peas (<i>Pisum</i> spp.) only
PRODUCT RATE (mL/ha)	500
APPLICATION TIMING	Make first application at the R1 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.
DISEASE	Cercospora Leaf Spot (<i>Cercospora kikuchii</i>)
CROPS	Soybeans only
PRODUCT RATE (mL/ha)	500
APPLICATION TIMING	Make first application at the R1 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.
DISEASES	<p>Ascochyta blight (<i>Ascochyta</i> spp.)</p> <p>Powdery Mildew (<i>Microsphaera diffusa</i>, <i>Erysiphe pisi</i>, <i>E. polygoni</i>)</p>

CROPS	Succulent shelled peas varieties
PRODUCT RATE (mL/ha)	500
APPLICATION TIMING	For Ascochyta blight the first application must be applied before disease is established and no later than the onset of flowering. A second application can be made 10-14 days after the first application, when disease pressure is severe or when agronomic or weather conditions are conducive to disease development or movement. For powdery mildew make first application at the R1 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.
DISEASES	Ascochyta blight (<i>Ascochyta</i> spp.) Mycosphaerella blight (<i>Mycosphaerella pinodes</i>) Anthracnose (<i>Colletotrichum</i> spp.) Sclerotinia (<i>Sclerotinia sclerotiorum</i>) – suppression only
CROPS	Crop Subgroup 6C: Dried shelled pea and bean (except soybean) - Any dried cultivar of bean (<i>Phaseolus</i> spp.); and dried cultivar of pea (<i>Pisum</i> spp.); dried cultivars of bean (<i>Lupinus</i> spp.) (includes grain lupin, sweet lupin, white lupin, and sweet white lupin); bean (<i>Vigna</i> spp.) (includes adzuki beans, mung bean, rice bean, southern bean, urd bean); bean (<i>Phaseolus</i> spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); broad bean (dry, fava bean); chickpea; guar; lablab bean; lentil; pea (<i>Pisum</i> spp.) (includes field pea); and pigeon pea.
PRODUCT RATE (mL/ha)	500
APPLICATION TIMING	The first application must be applied before disease is established and no later than the onset of flowering. A second application can be made 10-14 days after the first application, when disease pressure is severe or when agronomic or weather conditions are conducive to disease development or movement.
Restrictions: <ol style="list-style-type: none"> DO NOT apply within 15 days of harvest (PHI 15 days). APPLY A MAXIMUM OF 2 APPLICATIONS OF ELATUS A FUNGICIDE PER SEASON. Not all members of the legume vegetable group have been tested for efficacy and phytotoxicity at the recommended label rates, and should be used at the discretion of the user. Do not feed dried pea vines to livestock. 	

12.0 MINOR USES

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Syngenta Canada Inc. under the User Requested Minor Use Label Expansion program. For these uses, Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

12.1 Sugarbeets

SUGARBEETS	
DISEASES CONTROLLED	Rhizoctonia root and crown rot (<i>Rhizoctonia solani</i>)
PRODUCT RATE	0.5 – 1.1 L product/ha (4-6 mL/100 m of row for in-furrow) Apply in 50 - 100 L water/ha.
NOTES	Apply once in-furrow at seeding or a banded application over the row soon after emergence but before the 6th leaf stage.
Restrictions: <ol style="list-style-type: none"> 1. The pre-harvest interval (PHI) is 100 days. 2. DO NOT apply by air. 3. Do not apply more than one application per year. 	

13.0 RESISTANCE-MANAGEMENT RECOMMENDATIONS

For Resistance management, please note that ELATUS A Fungicide contains a Group 11 fungicide. Any fungal population may contain individuals naturally resistant to ELATUS A Fungicide and other Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of ELATUS A Fungicide or other Group 11 fungicides with different groups that control the same pathogens.

Do not exceed the total number of applications of ELATUS A Fungicide per season per crop as stated in Table 2.

Do not apply sequential treatments of ELATUS A Fungicide, or other fungicides in the same Fungicide Group, in a season. Do not apply at rates lower than recommended on the label.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for resistance development. Notify Syngenta Canada Inc. if reduced sensitivity of the pathogen to ELATUS A Fungicide is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, to which pathogen resistance has not developed.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

For further information and to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

Application Limitation and Preharvest Interval (PHI)

Crop	PHI	Maximum number of applications per year
Legume Vegetables	15 days	2
Chickpeas	15 days	2
Potato	N/A	1
Sugarbeets	100 days	1

ELATUS® is a trademark of a Syngenta Group Company.

GROUP

7

FUNGICIDE

ELATUS® B Fungicide
SOLATENOL® technology

FUNGICIDE

EMULSIFIABLE CONCENTRATE

COMMERCIAL - AGRICULTURAL

For broad-spectrum disease control or suppression of diseases in labelled crops.

ACTIVE INGREDIENT:

Benzovindiflupyr..... 100 g/L

READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**POISON**
DANGER: CORROSIVE TO EYES & SKINREGISTRATION NO: **31977**
PEST CONTROL PRODUCTS ACTNET CONTENTS: **0.5 – 1000 L****Syngenta Canada Inc.**
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

Label

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically. Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Keep unused product in original container tightly closed, locked up and away from food and feed.

CORROSIVE to the eye. **DO NOT** get in eyes. Causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if inhaled. Avoid inhaling/breathing sprays. Severely irritating to the skin. **DO NOT** get on skin.

DO NOT re-enter treated fields for 12 hours after application (12 hr REI).

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield) during mixing, loading, application, clean up and repair. Protective eyewear, chemical resistant gloves and coveralls are not required during application within a closed cab and/or cockpit.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Benzovindiflupyr is persistent and may carryover. It is recommended that this product not be used in areas treated with any products containing benzovindiflupyr during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

7.0 STORAGE

Store this product away from food or feed. Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area out of the reach of children and animals. This product does not require heated storage.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill,

and for cleanup of spills.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

ELATUS® and SOLATENOL® are trademarks of a Syngenta Group Company.

GROUP

7

FUNGICIDE

ELATUS® B Fungicide
SOLATENOL® technology

FUNGICIDE

EMULSIFIABLE CONCENTRATE

COMMERCIAL - AGRICULTURAL

For broad-spectrum disease control or suppression of diseases in labelled crops.

ACTIVE INGREDIENT:

Benzovindiflupyr..... 100 g/L

READ THE LABEL AND THIS PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**POISON**
DANGER: CORROSIVE TO EYES & SKINREGISTRATION NO: **31977**
PEST CONTROL PRODUCTS ACT**Syngenta Canada Inc.**
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

Pamphlet

Table of Contents

	<u>Section Number</u>
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE)	5.0
ENVIRONMENTAL PRECAUTIONS	6.0
STORAGE	7.0
DISPOSAL	8.0
PRODUCT INFORMATION	9.0
General Use Precautions	9.1
DIRECTIONS FOR USE	10.0
General Information	10.1
Ground Application	10.2
Mixing Instructions	10.2.1
Spraying Instructions	10.2.2
Equipment Clean-Up	10.2.3
Aerial Application	10.3
Operator Precautions	10.3.1
Pilot Precautions	10.3.2
Mixing Instructions	10.3.3
Spraying Instructions	10.3.4
Equipment Clean-Up	10.3.5
Product Specific Precautions	10.3.6
Spray Buffer Zones	10.4
Rotational Crop Restrictions	10.5
CROP USE DIRECTIONS	11.0
Potatoes – In Furrow Application	11.1
Sugar Beets	11.2
Dried Shelled Pea and Bean (except soybean) (Crop Subgroup 6C)	11.3
Soybeans	11.4
RESISTANCE MANAGEMENT RECOMMENDATIONS	12.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically. Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Keep unused product in original container tightly closed, locked up and away from food and feed.

CORROSIVE to the eye. **DO NOT** get in eyes. Causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if inhaled. Avoid inhaling/breathing sprays. Severely irritating to the skin. **DO NOT** get on skin.

DO NOT re-enter treated fields for 12 hours after application (12 hr REI).

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield) during mixing, loading, application, clean up and repair. Protective eyewear, chemical resistant gloves and coveralls are not required during application within a closed cab and/or cockpit.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Benzovindiflupyr is persistent and may carryover. It is recommended that this product not be used in areas treated with any products containing benzovindiflupyr during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

7.0 STORAGE

Store this product away from food or feed. Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area out of the reach of children and animals. This product does not require heated storage.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill,

and for cleanup of spills.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

9.0 PRODUCT INFORMATION

ELATUS® B is a broad-spectrum product and is recommended for the control or suppression of many important plant diseases. ELATUS B provides disease control of many rust, leaf spots and powdery mildews. ELATUS B is primarily applied as a foliar spray and can be used in block, or alternating spray programs with other crop protection products. All applications should be made according to the use directions listed in the label.

9.1 General Use Precautions

Do not apply ELATUS B through any type of irrigation equipment.

DO NOT USE IN GREENHOUSES (ALONE OR IN TANK MIXTURES).

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Apply the higher rate and shorter interval under conditions of high disease pressures.

Unless indicated otherwise in the use directions for specific crops, make no more than two sequential application of ELATUS B before alternating with a non-Group 7 fungicide registered for the same disease.

Not all members of the crop groups have been tested for efficacy and phytotoxicity at the recommended label rates, and should be used at the discretion of the user.

10.0 DIRECTIONS FOR USE

10.1 General Information

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Thorough coverage is necessary to provide good disease control. Applications using nozzles, sufficient water volume and pressure to provide thorough and uniform coverage generally provide the most effective disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

For more information on spray equipment and calibration, consult sprayer manufacturers and provincial recommendations.

Do not apply through any ultra-low volume (ULV) spray system.

It is important to check the physical compatibility of tank mixed pesticide products in a small volume prior to filling the sprayer. Check the compatibility of tank mixes containing A15457 Fungicide using a jar test with proportionate amounts of mix partners, and water, before mixing in the spray tank.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

All directions for use, crops/sites, use rates, dilution rates, precautions, and limitations which appear on the tank-mix product label must be observed. The label dosage for the tank-mix partner is not to be exceeded, and the most restrictive label precautions and limitations are to be followed.

10.2 Ground Application

- Apply in a minimum of 150 L of water per hectare, unless specified otherwise.
- Do not apply through any ultra-low volume (ULV) spray system.
- Thorough coverage is necessary to provide good disease control.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

10.2.1 Mixing Instructions

Prepare only the amount of spray mixture that is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

- ELATUS B is an emulsifiable concentrate (EC) formulation.
- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

ELATUS B Alone (No Tank Mix)

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add ELATUS B to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after ELATUS B has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

ELATUS B + Tank Mixtures: ELATUS B is usually compatible with all tank-mix partners listed on this label. To determine the physical compatibility of ELATUS B with other products, use a jar test. Using a jar, add the products to water in the same ratio and order as planned in the spray tank. Add wettable powders and water dispersible granular products first, then liquid flowables, then emulsifiable concentrates and solutions last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same order of addition required ingredients to the spray tank.

Tank Mixtures: All directions for use, crops/sites, use rates, dilution rates, precautions, and limitations which appear on the tank-mix product label must be observed. The label dosage for the tank-mix partner is not to be exceeded, and the most restrictive label precautions and limitations are to be followed.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by Syngenta Canada Inc.

1. Ensure that the sprayer interior is clean, then fill the spray tank with $\frac{1}{2}$ the required amount of water and engage gentle agitation. Good agitation is indicated by a rippling or rolling action on the surface of the water.
2. Add any WG or DF formulation mix partners and agitate to ensure complete mixing.
3. Add any additional SC formulation mix partners and agitate to ensure complete mixing.
4. Add ELATUS B and agitate to ensure complete mixing.

5. Add any other EC formulation mix partners and agitate to ensure complete mixing.
6. Fill the tank to $\frac{3}{4}$ the required amount of water.
7. Add any solution (SN or SL) formulation mix partners and agitate to ensure complete mixing.
8. Finish filling the sprayer with water, maintaining good agitation.
9. After any break in spraying operations, agitate thoroughly before spraying again.
10. Spray the pesticide suspension the same day as mixing.
11. **DO NOT** mix, load or clean spray equipment where there is a potential to contaminate wells or aquatic systems.

When using chemical handling equipment to fill the sprayer, the following additional recommendations apply:

- WG and DF formulations are preferentially batch mixed.
- SC, SN, and SL formulations may be inducted or batch mixed.
- EC formulations are preferentially batch mixed.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and ELATUS B to the spray tank.
- Allow ELATUS B to completely disperse.
- Spray the mixture with the agitator running.

10.2.2 Spraying Instructions

1. Water Volume: Apply in a minimum spray volume of 150 L/ha OR the volume given in the crop and pest specific instructions tabulated below, whichever is LARGER.
2. Sprayer Agitation: Use a jet agitator or liquid sparge tub which recirculates 10% or more of the tank per minute. **DO NOT** use an air sparger.
3. Pump: Screens should be used to protect the pump and prevent clogging. Use 16 mesh or *coarser* screens on the suction side of the pump. **DO NOT** place a screen in the recirculation line. Use 50 mesh or *coarser* screens between the pump and boom.
4. Spray Nozzles: 80° or 110° drift reducing flat fan (e.g. those with a pre-orifice or turbulence chamber) or air induction nozzles are recommended. Use 50 mesh nozzle screens. **DO NOT** use flood type nozzles, controlled droplet application equipment, spray foils or hollow cone nozzles.
5. Pressure: As recommended by the nozzle manufacturer to achieve ASAE medium sized droplets.
6. Apply at uniform speed and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to avoid potential crop injury from over application.

10.2.3 Equipment Clean-Up

Before Spraying:

Prior to using ELATUS B, ensure that the spray tank, chem handling equipment, lines and filter are thoroughly cleaned.

After Spraying:

Thoroughly clean application equipment immediately after spraying. **DO NOT** let ELATUS B residue to dry within application equipment.

When using tank mixes, consult the tank-mix partner label for additional clean-up instructions.

The following recommendations are provided:

1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **DO NOT** clean application equipment near desirable vegetation, wells or other water sources.
2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial regulations.

10.3 Aerial Application

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions, and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. When no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

10.3.1 Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

10.3.2 Pilot Precautions

Read and understand the entire label before using this product. Apply only when meteorological

conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides. Light winds at altitude of 3-16 km/h are preferred, with 3-9 km/h considered optimal. Ensure uniform application and a uniform spray with minimum potential for drift. To avoid streaked, uneven or overlapped application, use appropriate marking technology. GPS based marking is preferred.

10.3.3 Mixing Instructions

Mixing this product directly in the aircraft hopper **IS NOT** recommended. The use of chemical handling or managing equipment to load the hopper **IS** recommended. This product **MAY BE** inducted into a hopper which is prefilled with water or when the product and water are mixed prior to entering the hopper. This product **MAY BE** batch mixed and pumped into the hopper. In all cases the chemical handling equipment and hopper interior must be clean prior to use.

NOTE: WG and DF formulations are preferentially batch mixed.

NOTE: SC, SN, and SL formulations may be inducted or batch mixed.

NOTE: EC formulations are preferentially batch mixed.

It is **NOT** recommended to combine solid (WG or DF) formulations with liquid tank mix partners within a single batch. Batch mix WG or DF formulations first, pump into the hopper, and then add liquid tank mix partners by induction or as an additional batch mix. When tank mixing multiple products, follow the mixing order outlined below:

- Pump water into the hopper to at least $\frac{1}{4}$ to $\frac{1}{2}$ of the desired spray volume. Engage hopper circulation, if possible.
- Thoroughly batch mix any WG or DF formulation mix partners and agitate to ensure complete mixing. Pump into the hopper
- Induct or thoroughly batch mix any SE or SC formulations.
- Thoroughly batch mix A15457 Fungicide and any EC formulation mix partners. EC formulations may be added to the batch from Step 3, if desired.
- Induct or thoroughly batch mix any solution (SN or SL) formulation mix partners. SN/SL formulations may be added to the batch from Step 3, if desired.
- Pump batch mixed SC, EC, and/or SN/SL products into the hopper.
- Finish filling the hopper with water.
- If it was not possible to engage hopper agitation in Step 1, do so as soon as possible once airborne.
- Spray the pesticide suspension the same day as mixing.
- **DO NOT** mix, load or clean equipment where there is a potential to contaminate wells or aquatic systems.

10.3.4 Spraying Instructions

1. Water Volume: Apply in a minimum spray volume of 45 L/ha OR the volume given in the crop and pest specific instructions tabulated below, whichever is LARGER.
2. Spray Nozzles: Use only ASAE medium or coarse nozzles rated as delivering droplets of volume median diameter of 300 microns or greater.
3. Pressure: As recommended by the nozzle manufacturer to achieve ASAE medium sized droplets.

4. Ensure hopper agitation is engaged whenever possible during flight.

10.3.5 Equipment Clean-Up

Before Spraying:

Prior to using ELATUS B, ensure that the hopper, chem handling equipment, lines and filter are thoroughly cleaned.

After Spraying:

Thoroughly clean application equipment immediately after spraying. **DO NOT** let ELATUS B residue to dry within application equipment.

When using tank mixes, consult the tank-mix partner label for additional clean-up instructions.

The following recommendations are provided:

1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **DO NOT** clean application equipment near desirable vegetation, wells or other water sources.
2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial regulations.

10.3.6 Product Specific Precautions

Follow recommendations provided by local disease monitoring services or provincial spray calendars regarding the appropriate application timing for protectant fungicides in your area. Cultural practices such as canopy management and removal of overwintered plant debris should be integrated with the use of fungicides to reduce disease incidence. Use the higher rate, and the shorter application interval under conditions of heavy infection pressure, on highly susceptible varieties, or when environmental conditions are favourable for disease development.

ELATUS B treatments should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, plant residue management, crop rotation, and proper timing and placement of fertilizer and irrigation. Fungicide applications should begin prior to disease infection and continue throughout the season following a resistance management strategy (see **Resistance Management Recommendations**).

Read and understand the entire label before opening this product. If you have questions, call Syngenta Canada Inc. at 1-877-964-3682 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum spray volume of 45 litres per hectare.

10.4 Spray Buffer Zones

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

Use of a low-clearance hooded or shielded sprayers that prevent contact with crop, fruit, or foliage, soil drench or soil incorporation do not require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Potatoes		15	2	1	1	1
Field sprayer	Dried shelled pea and bean subgroup (legumes), soybeans		10	1	1	1	1
Aerial	Dried shelled pea and bean subgroup (legumes), soybeans	Fixed wing	675	15	5	1	15
		Rotary wing	325	10	2	1	10

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

10.5 Rotational Crop Restrictions

Please see the following table for the crop rotational restrictions:

Rotational Crops	Planting Time From Last ELATUS B Application
Potatoes and tuberous and corm vegetable subgroup (CG 1C) Dried shelled pea and bean subgroup (CG 6C) Soybeans Fruiting vegetables crop group (CG 8-09) Cucurbit vegetables crop group (CG 9) Cereals (wheat, barley, oats, rye, triticale) Grasses grown for seed (timothy, annual and perennial ryegrass) Corn (field, sweet, pop and specialty) Rapeseed subgroup (CG 20A) Onions (Crop Subgroups 3-07A and 3-07B) Sugar Beets	0 days
All other crops Intended for Food and Feed	6 months/180 days

11.0 CROP USE DIRECTIONS

Apply the higher rate and shorter interval under conditions of high disease pressures.

Diseases are controlled unless otherwise indicated as suppressed.

11.1 Potatoes – In Furrow Application

Application Timing/Instructions: Make an in-furrow application in a water volume of 50-150 L per hectare at planting. Mount the spray nozzle so the spray is directed into the furrow as a 15-20 cm band just before the seed is covered. Apply the spray in a narrow band over the seed piece. Do not apply by air.

Specific Use Restrictions:

Do not apply more than 100 g a.i./ha of foliar benzovindiflupyr-containing products on potato crops per season if an in-furrow application was made.

Diseases suppressed at 500-750 mL product/ha. See Rate Conversion Chart – Potatoes for rates per 100 m row	Stem and stolon canker and black scurf (<i>Rhizoctonia solani</i>)
Diseases suppressed at 750 mL product/ha. See Rate Conversion Chart – Potatoes for rates per 100 m row	Verticillium wilt (<i>Verticillium dahliae</i>)
Diseases suppressed at 500-750 mL product/ha. See Rate Conversion Chart – Potatoes for rates per 100 m row.	Silver scurf (<i>Helminthosporium solani</i>)

RATE CONVERSION CHART - POTATOES

Rate mL product /ha	Rate in mL product /100 m row for specified row spacing									
	Row Spacing (cm)									
	75	80	85	90	95	100	105	110	115	120
500	3.8	4.0	4.3	4.5	4.8	5.0	5.3	5.5	5.8	6.0
750	5.6	6.0	6.4	6.8	7.1	7.5	7.9	8.3	8.6	9.0

11.2 Sugar Beets

Application Timing/Instructions: Apply once in-furrow at seeding with a minimum water volume of 50 L/ha or a banded application with a minimum water volume of 100 L/ha over the row soon after emergence but before the 6th leaf stage. If cool soil conditions are expected after planting which could result in an extended period of plant emergence, do not apply ELATUS B in-furrow.

Diseases suppressed at 500-750 mL product/ha.	Rhizoctonia root and crown rot (<i>Rhizoctonia solani</i>)
--	---

11.3 Dried Shelled Pea and Bean (except Soybean) (Crop Subgroup 6C)

Crops	Dried shelled pea and beans (Crop Subgroup 6C) Any dried cultivar of bean (<i>Phaseolus</i> spp.); and dried cultivar of pea (<i>Pisum</i> spp.): dried cultivars of bean (<i>Lupinus</i> spp.) (includes grain lupin, sweet lupin, white lupin, and sweet white lupin); (<i>Vigna</i> spp.) (includes adzuki beans); (<i>Phaseolus</i> spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean, mung bean, rice bean, southern bean, urd bean); broad bean (dry, faba bean); chickpea; guar; lablab bean; lentil; pea (<i>Pisum</i> spp.) (includes field pea); and pigeon pea
Diseases Controlled	Ascochyta blight (<i>Ascochyta</i> spp.) Asian soybean rust (<i>Phakopsora pachyrhizi</i>) Anthracnose (<i>Colletotrichum</i> spp.)
Use rate (mL product/ha)	500-750
Application Timing/Instructions	Make the first application prior to disease development. A second application can be made 14 days after the first application, when disease pressure is severe or when agronomic or weather conditions are conducive to disease development or movement. If disease pressure is high, use the highest rate and shortest interval. For best results, sufficient water volume must be used to provide thorough coverage. A minimum of 100-200 L/ha for ground applications is recommended. For aerial applications, a minimum of 45 L/ha of water is recommended.
Specific Use Restrictions: <ol style="list-style-type: none"> 1. Do not apply more than 1.5 L/ha/season of ELATUS B. 2. Do not apply within 15 days of harvest for dry shelled peas and beans (15-day PHI). 3. Dry pea hay and vines may be fed or harvested 15 days (15-day PHI) after last application. 4. Do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program. 5. Not all members of the legume vegetable subgroup 6C have been tested for efficacy and phytotoxicity at the recommended label rates, and should be used at the discretion of the user. 	

11.4 Soybeans

Crops	Soybeans
Diseases Controlled	Septoria brown spot (<i>Septoria glycines</i>) Frogeye leaf spot (<i>Cercospora sojina</i>) (suppression) Asian soybean rust (<i>Phakopsora pachyrhizi</i>) Pod and stem blight (<i>Diaporthe phaseolorum</i>) (suppression)
Use rate (mL product/ha)	500-750
Application Timing/Instructions	Make the first application prior to disease establishment. A second application can be made 7-14 days after the first application, when disease pressure is severe or when agronomic or weather conditions are conducive to disease development or movement. If disease pressure is high, use the highest rate and shortest interval. For best results, sufficient water volume must be used to provide thorough coverage. A minimum of 100-200 L/ha for ground applications is recommended. For aerial applications, a minimum of 45 L/ha of water is recommended.
Specific Use Restrictions: <ol style="list-style-type: none"> 1. Do not apply more than 1.5 L/ha/season of ELATUS B. 2. Do not apply within 14 days of harvest for soybean grain (14-day PHI). 3. Soybean forage and hay may be fed or harvested 1 day (1-day PHI) after last application. 4. Do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program.	

12.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, ELATUS B contains a Group 7 fungicide in the succinyl dehydrogenase inhibitor (SDHI) class. Any fungal population may contain individuals naturally resistant to benzovindiflupyr and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of benzovindiflupyr or other Group 7 fungicides with different groups that control the same pathogens.

Apply no more than 2 sequential applications unless otherwise stated in the crop section.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for sign of resistance development.

If disease continues to progress after treatment with this product, DO NOT increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, to which pathogen resistance has not developed.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-877-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

Application Limitations and Pre-harvest Intervals

Crop*	Maximum Product Rate/ha/application (mL/ha)	Maximum total (L/ha/Season)	Pre-Harvest Interval (PHI) (days)	Minimum Re-treatment Interval (days)
Dried Shelled Pea and Bean (CG 6C)	750	1.5	15	14
Pea hay and vines	750	1.5	15	14
Soybeans	750	1.5	14	7
Soybean forage and hay	750	1.5	1	7
Potatoes	750 (soil)	0.75 (soil)	N/A	N/A
Sugarbeets	750 (soil)	0.75 (soil)	N/A	N/A

*For specific crops in a group and use directions, refer to the Specific Directions For Use

ELATUS® and SOLATENOL® are of a Syngenta Group Company.